

# Constraining branon dark matter from observations of dwarf spheroidal galaxies with the MAGIC Telescopes

Tjark Miener (tmiener@ucm.es), Daniel Nieto, Viviana Gammaldi, Javier Rico



**Grupo de Altas Energías - UCM**

VII Meeting on Fundamental Cosmology (11/09/2019)

# Outline

## Constraining branon dark matter

Introduction

Branon dark matter

gLike tool

Results

# Introduction





Figure: Dark matter (DM) self-annihilation. [Fermi-LAT]

# Introduction



Expected gamma-ray flux produced by DM annihilation:


$$\frac{d\Phi(\Delta\Omega, E_{\gamma})}{dE_{\gamma}} = \underbrace{\mathcal{J}(\Delta\Omega)}_{\text{Astrophysics}} \cdot \underbrace{\frac{1}{4\pi} \frac{\langle\sigma_{ann}v\rangle}{2m_{\chi}^2} \sum_i \text{BR}_i \frac{dN_{\gamma}^i}{dE_{\gamma}}}_{\text{Particle physics}}$$

$$\mathcal{J}(\Delta\Omega) = \int_{\Delta\Omega} d\Omega' \int_{\text{l.o.s.}} dl \rho^2(l, \Omega')$$

Figure: Dark matter (DM) self-annihilation. [Fermi-LAT]

# MAGIC telescope & Segue 1 observations

- ▶ Currently operating imaging atmospheric Cherenkov telescope
- ▶ Roque de los Muchachos Observatory on La Palma at about 2200m above sea level
- ▶ Two telescopes with 17m diameter reflecting surfaces placed at a distance of 85m
- ▶ Sensitive to VHE gamma-rays (between  $\sim 50$  GeV and  $\sim 50$  TeV) [MAGIC]
- ▶ Segue 1 data set is almost 160 hours of good-quality data and was taken under four different experimental conditions



# Dark matter particle zoo






Figure: Summary of the DM particle zoo candidates. [J. Conrad]

# Brane World Theory

- ▶ Dark matter cannot be made of any of the known particles of the Standard Model (SM).
- ▶ Our work focuses on Brane World Theory as a prospective framework for DM candidates.
- ▶ Branons are new degrees of freedom that appear in flexible brane-world models corresponding to brane fluctuations.
- ▶ Branons behave as Weakly Interacting Massive Particles (WIMPs), that are one of the most favored candidates for DM.


# Branon dark matter

$$\frac{d\Phi(\Delta\Omega, E_\gamma)}{dE_\gamma} = \underbrace{\mathcal{J}(\Delta\Omega)}_{\text{Astrophysics}} \cdot \underbrace{\frac{1}{4\pi} \frac{\langle\sigma_{ann}v\rangle}{2m_\chi^2} \sum_i \text{BR}_i}_{\text{Particle physics}} \frac{dN_\gamma^i}{dE_\gamma}$$






# Annihilation photon yield for $m_\chi = 1$ TeV



# Branon annihilation photon yields



- ▶ gLike is a code framework for the numerical maximization of joint likelihood functions.
- ▶ gLike can estimate the dark matter annihilation cross-section combining observations of dark matter targets by different ground-based gamma-ray telescopes, satellite gamma-ray detectors, neutrino telescopes, ...
- ▶ Open source on GitHub: <https://github.com/javierrico/gLike>

## Results - First limit to branon dark matter





Figure: Limit to branon dark matter annihilation cross-section from MAGIC observations of Segue 1.

## Summary & Outlook

- ▶ We analyzed the MAGIC Segue 1 high-level data set, which is the deepest IACT observational campaign on any dwarf galaxy.
- ▶ We performed a full joint likelihood analysis from four different instrument conditions.
- ▶ We modified gLike to include the branon dark matter model in our analysis.
- ▶ We would like to combine more observations from other dark matter targets from different instruments to improve our branon limits.



Gracias por su atención!



Back up

# Segue1 observational campaign with MAGIC

	Sample A	Sample B1	Sample B2	Sample C
Readout	DRS2	DRS4	DRS4	DRS4
MAGIC-I camera	old	old	old	new
Obs. period	Jan–May 2011	Jan–Feb 2012	Mar–May 2012	Nov 2012–Feb 2013
Obs. time [h]	64	24.28	59.77	55.05
$Zd$ range [deg]	13–33.7	13–32.5	13–35.7	13–37
$Az$ range [deg]	104.8–250.2	120.2–252.0	115.4–257.2	103.8–259.4
Wobble around	dummy	dummy	dummy	Segue 1
Wobble offset [deg]	0.29	0.29	0.29	0.40
W1 $t_{\text{eff}}$ [h]	22.66	6.07	25.02	23.71
W2 $t_{\text{eff}}$ [h]	24.35	6.20	26.11	23.80
$t_{\text{eff}}$ [h]	47.00	12.26	51.13	47.51
<b>Total <math>t_{\text{eff}}</math> [h]</b>				<b>157.9</b>

**Table 1:** Basic details of the Segue 1 observational campaign with MAGIC. Refer to the main text for additional explanations.

Figure: [MAGIC]



## Profile likelihood

- ▶ We are interested in the parameter  $g = (g_1, \dots, g_k)$ . (For DM searches  $g$  is  $\langle \sigma v \rangle$ )
- ▶ The model describing our data depends on  $g$  and additional nuisance parameters  $h = (h_1, \dots, h_l)$ .
- ▶ The full likelihood function is given by

$$\mathcal{L}(g; h|X) = \prod_{i=1}^n f(X_i|g; h),$$

where  $X = (X_1, \dots, X_n)$  are  $n$  independent observations and  $f(X|g; h)$  is the probability density function PDF.

# Joint likelihood

- ▶ Combining likelihood functions for different targets:

$$\mathcal{L}(\langle \sigma v \rangle; \nu | \mathbf{X}) = \prod_{i=1}^{N_{\text{target}}} \mathcal{L}_i(\langle \sigma v \rangle; J_i, \mu_i | X_i) \cdot \mathcal{J}(J_i | J_{\text{obs},i}, \sigma_i)$$

- ▶ Combining likelihood functions (of a particular target) for different experiments:

$$\mathcal{L}_i(\langle \sigma v \rangle; J_i, \mu_i | X_i) = \prod_{j=1}^{N_{\text{instrument}}} \mathcal{L}_{ij}(\langle \sigma v \rangle; J_i, \mu_{ij} | X_{ij})$$