Dark matter annual modulation and ANAIS-112 results:

testing the DAMA/LIBRA positive signal

María Martínez, ARAID & U. Zaragoza

CIEMAT, October 29, 2020

OUTLINE

- Intro: Dark matter annual modulation
- DAMA/LIBRA positive signal
- Current NaI(Tl) experiments
- ANAIS-112
  - Experimental set-up
  - Detector performance
  - Results on annual modulation
  - ANAIS-112 sensitivity
- Summary

Intro: DARK MATTER ANNUAL MODULATION



Evidence of Dark Matter



Zwicky (1933)













DM dominates all the structures of the Universe!







DM candidates

What we know:

- lifetime > > age of Universe
- No (or very very small) interaction with light
- Non baryonic
- Cold (or Warm) (Moving non-relativistically when galaxies started to form)
- Beyond the Standard Model



WIMPs

Weakly Interacting Massive Particles (WIMPs) very well motivated

- The relic abundance determined from the freeze-out mechanism matches the measured DM density for reasonable ranges of weak-scale annihilation cross section
- WIMPs arise spontaneously in many extensions of the Standard Model such as SUSY



Jungman *et al* hep-ph/9506380

Abundance of a thermal relic

$$\sim \frac{0.1\ pb}{<\sigma_A\ v/c>}$$



WIMP direct detection

Dark Matter Halo

Density $\rho_0 \sim 0.4$ GeV / cm^3

Milky Way

Earth

V ~ 200 km/s

$$\phi^{WIMP}{}_{earth} \sim 10^8 - 10^{10}\ s^{-1}\ m^{-2}$$



WIMP direct detection

WIMPs scatter elastically off nuclei



WIMP direct detection

WIMPs scatter elastically off nuclei



Expected rate @ Earth:

MODEL DEPENDENT!

$$\frac{dR}{dE_R} = \frac{\rho_0 M_{Det}}{2 m_W m_{WN}^2} \sigma_{WN} \int_{vmin}^{vmax} \frac{f(v)}{v} dv^3$$

Wimp model

Halo model



Naïve approximation: **SHM**

Isothermal sphere

Maxwellian distribution

v_o= 220 km/s

Truncated at v_{esc} ~ 530 km/s



DM Annual modulation

$$v_0 = 220\ km/s$$

$$\frac{dR}{dE_R} = \frac{\rho_0 M_{Det}}{2 m_W m_{WN}^2} \sigma_{WN} \int_{vmin}^{vmax} \frac{f(v)}{v} dv^3$$



DM Annual modulation



$$\frac{dR}{dE_R} = \frac{\rho_0 M_{Det}}{2 m_W m_{WN}^2} \sigma_{WN} \int_{vmin}^{vmax} \frac{f(v,t)}{v} dv^3$$



DM Annual modulation



Due to the Earth revolution around the Sun, the relative speed Earth-halo is cosine-like with 1 year periodicity and small amplitude ($\sim 7\%$), and that implies **a modulation in the expected rate:**

$$\frac{dR}{dE}(E,t) \approx S_0(E) + S_m(E)\cos\omega(t-t_0)$$

Where

$$S_m(E) = \frac{1}{2}\left(\frac{dR}{dE}(E,t_0) - \frac{dR}{dE}(E,t_0+182)\right)$$







A distinctive signal hard to mimic by background

- ✓ **Cosine behaviour**
- ✓ **1 year period**
- ✓ **Maximum around June 2nd**
- ✓ **Weak effect** $\left(\frac{S_m}{S_0} \sim 0.01 - 0.1\right)$
- ✓ **Only noticeable at low energy**
- ✓ **Phase reversal at low E**

DAMA: an observatory for rare processes @LNGS

http://people.roma2.infn.it/dama

DAMA/NaI & DAMA/LIBRA (phase 1)

DAMA / NaI (1995-2002)



- 10 × 9.7 kg NaI(Tl) (3x3 matrix)
- 7 annual cycles
- Exposure : 0.29 ton × y



Riv. Nuov. Cim., 26 (2003) 1

The data support the presence of a modulation (1 y period, phase on May 20th) at **6.3σ CL**

DAMA/NaI & DAMA/LIBRA (phase 1)

DAMA / NaI (1995-2002)



- 10 × 9.7 kg NaI(Tl) (3x3 matrix)
- 7 annual cycles
- Exposure : 0.29 ton × y



Riv. Nuov. Cim., 26 (2003) 1

The data support the presence of a modulation (1 y period, phase on May 20th) at **6.3σ CL**

DAMA / LIBRA (2003-2010)



- 25 × 9.7 kg NaI(Tl) (5x5 matrix)
- 7 annual cycles
- Exposure : 1.17 ton × y



Eur. Phys. J. C (2013) 73:2648

The data support the presence of a modulation (1 y period, phase on May 20th) at **9.2σ CL**

DAMA/NaI & DAMA/LIBRA (phase 1)

DAMA / NaI (1995-2002)



- 10 × 9.7 kg NaI(Tl) (3x3 matrix)
- 7 annual cycles
- Exposure : 0.29 ton × y

DAMA / LIBRA (2003-2010)



- 25 × 9.7 kg NaI(Tl) (5x5 matrix)
- 7 annual cycles
- Exposure : 1.17 ton × y





The signal satisfies all requirements for DM and **can be interpreted as a WIMP:**



Interpreting DAMA/LIBRA ph1 as WIMPs

DAMA clearly sees an annual modulation at 12.9σ



But **the parameter's region singled out by DAMA/LIBRA is excluded by most sensitive experiments**, even assuming more general halo/interaction models!

Other annual modulation searches

"A search for annual and diurnal rate modulations in the LUX experiment", Phys. Rev. D 98, 062005 (2018)



"Direct dark matter search by annual modulation with 2.7 years of XMASS-I data", Phys. Rev. D 97, 102006 (2018)



"Search for Electronic Recoil Event Rate Modulation with 4 Years of XENON100 Data" PRL118, 101101 (2017)



"Search for annual modulation in low-energy CDMS-II data", 1203.1309



None of them is compatible with DAMA/LIBRA

2011-2013 exciting times



Hint of annual modulation in one experiment

CoGeNT (2011-2014)



One PPC HPGe @ SOUDAN

330 g Ge, 3.4 years
No β/γ discrimination

(Sm 4-7 times larger than predicted in SHM)

Excess of events in WIMP region in two experiments:

CRESST (2011)



SuperCDMS-Si (2013)





ArXiv:1401.3295

2011-2013 exciting times



CoGeNT modulation too large, but can be in agreement with DAMA/LIBRA for some non-SHM haloes



But since 2013 … ☹

- CRESST-II (2014-2015) with an upgraded detector did not confirm the excess
- CoGeNT's signal significance decreased with time (3.4y by 2014) to below 2σ. Different interpretations among the collaboration.
- CDMS-II-Si signal very small. No more data

DAMA/LIBRA phase2 (2011-2018)

1805.10486, Nucl. Phys. At. Energy 19, 307 (2018)

all PMTs replaced with new ones of higher Q.E.



2 more points





- 6 anual cycles
- Exposure: 1.13 ton × y



The data of DAMA/LIBRA phase1+phase2 favor the presence of a modulation with proper features at **11.9σ CL** (2.17 ton × yr)

Interpretation of the 1 keV point



1804.01231, Baum, Freese, Kelso "Dark Matter implications of DAMA/LIBRA-phase2 results"

"the observed annual modulation signal is no longer well fitted by canonical (isospin conserving) spin-independent WIMP nucleon couplings"

Interpretation of the 1 keV point



1907.06405 , Bernabei et al. "Improved model-dependent corollary analyses after the first six annual cycles of DAMA/LIBRA-phase2"

"at present level of uncertainties the DAMA data, if interpreted in terms of DM particle inducing nuclear recoils through SI interaction, can account either for low and large DM particle mass and for a wide range of the ratio fn=fp, even including the "standard" case fn=fp = 1."



1804.01231, Baum, Freese, Kelso "Dark Matter implications of DAMA/LIBRA-phase2 results"

"the observed annual modulation signal is no longer well fitted by canonical (isospin conserving) spin-independent WIMP nucleon couplings"



"the purely SD scenarios are in good agreement with the DAMA results and can explain the different capability of detection among targets with different unpaired nucleon."

Interpreting DAMA/LIBRA ph1 as WIMPs

DAMA clearly sees an annual modulation at 12.9σ



Dark matter or systematics?

TO AVOID ANY MODEL DEPENDENCE, WE NEED A PROOF/DISPROOF WITH THE SAME TARGET

Model dependency

Expected rate @ Earth:

Wimp model

TARGET DEPENDENT!

$$\frac{dR}{dE_R} = \frac{\rho_0 M_{Det}}{2 m_W m_{WN}^2} \sigma_{WN} \int_{vmin}^{vmax} \frac{f(v)}{v} dv^3$$

Halo model

In order to compare experiments using different targets I have to assume:

- Only SI or SD coupling
- For SI: - Isospin conserving (couplings are identical for protons and neutrons)
  - A scaling law for the cross section

$$\sigma_{SI} \propto \frac{m_{WN}^2}{m_{Wn}^2} A^2 \; F^2 \sigma_{SI}^{nucleon}$$

- A model for the WIMPs velocity distribution, usually the standard Halo Model (Maxwellian distribution)



$$v_{min} = \sqrt{\frac{m_N E_{th}}{2\mu_{NW}^2}}$$

- A good knowledge of the NR quenching factor

Model dependency

Wimp model

TARGET DEPENDENT!

Expected rate @ Earth:

$$\frac{dR}{dE_R} = \frac{\rho_0 M_{Det}}{2 m_W m_{WN}^2} \sigma_{WN} \int_{vmin}^{vmax} \frac{f(v)}{v} dv^3$$

Halo model

In order to compare experiments using different targets I have to assume:

- Only SI or SD coupling
- For SI: - Isospin conserving (couplings are identical for protons and neutrons)
  - A scaling law for the cross section

$$\sigma_{SI} \propto \frac{m_{WN}^2}{m_{Wn}^2} A^2 \; F^2 \sigma_{SI}^{nucleon}$$

- A model for the WIMPs velocity distribution, usually the standard Halo Model (Maxwellian distribution)



$$v_{min} = \sqrt{\frac{m_N E_{th}}{2\mu_{NW}^2}}$$

- A good knowledge of the NR quenching factor

That is the only that **could** remain when using the same target

NR quenching factor

In a scintillator, an **ER** produces much more light than a **NR** of the same energy!



The spectra are calibrated with X/γ sources, so are given in keVee(*). In order to be interpreted as NR, $\boldsymbol{QF}$ **has to be measured** to correct the energy scale:

$$\boldsymbol{QF} = \frac{signal_{NR}/keV}{signal_{ER}/keV}$$

(*) keVee: electron-equivalent keV

The "true" NR energy is:



But I measure:



NR quenching factor measurements

DAMA uses (Spooner'94) NaI quenching factor measurements : $QF_{Na} = 0.3$ $QF_I = 0.09$
But recent measurements give lower values. Na quenching decreases when decreasing the energy.

Joo, et al. Astrop. Phys. 108 (2019) 50



Mod Amplitude in keVee WIMP 50 GeV

Iodine signal



Na signal





NR quenching factor measurements

DAMA uses (Spooner'94) NaI quenching factor measurements : $QF_{Na} = 0.3$ $QF_I = 0.09$
But recent measurements give lower values. Na quenching decreases when decreasing the energy.



Does the QF depend on the crystal?

- Impurities
- Tl level
- Crystal quality
- …

To answer this question,
Anais + Yale
QF measurements @ TUNL (Duke Univ.)
different NaI(Tl) crystals (ANAIS & COSINE)
in the same setup
Results soon!



CURRENT NaI(Tl) experiments

NaI experiments around the World



NaI experiments around the World



Why took so long?

$$\text{Sensitivity} \propto \sqrt{\frac{MT\epsilon}{B}}$$

Large mass

Stable conditions over years

High efficiency at very low energy

Very low radioactive background!!

- **The main contribution to the background comes from the crystal itself**
- Long effort of ANAIS team looking for ultra pure NaI(Tl), R&D with Alpha Spectra → crystals now used by ANAIS-112 and COSINE-100
- However, **up to the date, the quality of the DAMA crystals has not been reached by any group**



	K (ppb)	^{210}Pb (mBq/kg)
DAMA (Saint Gobain)	13	0.01-0.03
ANAIS/COSINE (Alpha Spectra)	18-44	0.7-3

COSINE-100

From Y. J. Ko @ TAUP2019



- Data-taking started in Sep 2016, Y2L (South Korea)
- 8 ultra low-background NaI(Tl) crystals with 106 kg in total (**but only ~60 kg usable for DM search**)
- Inside lead shielding and Liquid Scintillator tank to reject coincident events (^{40}K!)
- Muon veto & neutron monitoring



NEXT STEP: COSINE-200 (2020?)

Goal: Run 200 kg NaI(Tl) in the same set-up, with improved background (lower than DAMA/LIBRA)

Status: Power purification, crystal growing and handling facilities established, buy a factor 2 or more improvement in bkg is needed.

From G. Adhikari @ TAUP2019

COSINE-100 results

From Y. J. Ko @ TAUP2019

SET1 (59.5 days)

Background + WIMP signal is fit to data:



→ Model dependent exclusion of DAMA/LIBRA-phase1 (*)

- Spin Independent interaction
- Maxwellian velocity distribution



Nature 564 (2018) 83-86

(*) But not excluded in effective models (COSINE coll. & S. Scopel, JCAP 1906 (2019) 06, 048)

SET2 (1.7 y, 97.7 kg·year exposure)

Phys.Rev.Lett. 123 (2019) 031301





At 68.3% C.L., result is consistent with both a null hypothesis and DAMA/LIBRA's best fit value.



SABRE

From S. Copello @ TAUP2019



- Ultra-clean NaI(Tl) (Princeton)
- Two sites (LNGS/ Stawell)

Proof of Principle: one NaI crystal in LS vessel



LS Vessel ready at LNGS

First crystal (3.5 kg)
arrived @ LNGS on August 2019

	K (ppb)	^{210}Pb (mBq/kg)
SABRE	4	0.4
DAMA	13	0.01-0.03
ANAIS	18-44	0.7-3

PICOLON

From K.Fushimi @ TAUP2019

GOAL: Development of highly radiopure NaI(Tl) scintillator



Status of NaI(Tl) purification (~April 2019)

RI	Ingot26 (2015)	Ingot37 (2016)	Ingot71 (2018)	Ingot76 (2019)	Goal
Size	3"ϕX3"	4"ϕX3#	3"ϕX3"	5"ϕX4"(*)	5"ϕX5"
^{40}K (ppb)	2630	120	<20	<20	<20
^{232}Th (ppt)	0.4±0.5	3.7±0.5	1.7±0.2	--	<4
^{238}U (ppt)	4.7±0.3	5.9±0.3	9.7±0.8	4.4±0.2	<10
^{210}Pb (μBq/kg)	30±7	2300	1076	~560	<50
Method	Resin for Pb	I26+cation resin	double re-crystallization	Pb resin + double re-crystallization	

DAMA

13

10-33

COSINUS

First NaI detector with particle discrimination
(Two channel approach: HEAT and LIGHT)





With a moderate exposure of few O(100) kg-days , can confirm or rule-out a **nuclear recoil origin** of the DAMA/LIBRA dark matter claim

Schäffner, K. et al. J Low Temp Phys (2018). https://doi.org/10.1007/s10909-018-1967-3



Present threshold:
8.26 keV_{NR}
(Goal: 1 keV_{NR})

ANAIS-112 experiment

ANAIS-112

Annual Modulation with NaI Scintillators

J. Amaré, I. Coarasa, S. Cebrián, D. Cintas, E. García, M. Martínez, M.A. Oliván,
Y. Ortigoza, A. Ortiz de Solórzano, J. Puimedón, A. Salinas, M.L. Sarsa

GOAL:

Confirmation of DAMA-LIBRA modulation signal with the same target and technique
(but different experimental approach and environmental conditions)

THE DETECTOR:

3x3 matrix of 12.5 kg NaI(Tl) cylindrical modules = **112.5 kg** of active mass



WHERE:

At Canfranc Underground Laboratory, @ **SPAIN** (under **2450 m.w.e.**)



taking data since August 2017

ANAIS-112: experimental setup





- 9 NaI(Tl) cylindrical crystals (12.5 kg each) in 3x3 matrix
- Ultrapure NaI powder (Alpha Spectra Inc)
- Each coupled to two Hamamatsu R12669SEL2 PMT (QE ~40%)





ANAIS-112: Low energy calibration

Detectors equipped with a **Mylar window**!

Radon-free system for low energy calibration:

- **^{109}Cd sources** on flexible wires (radon-free)
- Energies: 11.9, 22.6 and 88.0 keV
- Simultaneous calibration of the nine modules
- Performed every two weeks





ANAIS-112: Muon Veto



In ANAIS we flag every muon that cross the shielding and set a (configurable) dead-time after every passage (DAMA/LIBRA has no muon veto)

The underground muon flux is annual-modulated!





DAMA reply:

- Modulation phase inconsistency
- Muons interacting directly in the detectors do not fulfill the DM requisites
- Not enough muon-induced fast neutrons to account for the signal



But still some open questions:

- (delayed) effect of muons in PMTs?
- slow phosphorescence in NaI?

ANAIS can test these hypotheses

ANAIS-112: Data acquisition system

- Individual PMT signals **digitized** and fully processed **(14 bits, 2 GS/s)**
- Trigger at phe level for each PMT signal
- AND coincidence in 200 ns window
- Redundant energy conversion by QDC
- Trigger in OR mode among modules
- Electronics at air-conditioned-room to decouple from temperature fluctuations
- Muon detection system: tag every muon event to offline processing









ANAIS-112: Slow control

- Monitoring **environmental parameters** since the start of DM run
  - Monitoring:
    Rn content, humidity, pressure, different temperatures, N_2 flux, PMT HV, muon rate, …
    Data saved every few minutes and alarm messages implemented
  - Stability checks:
    gain, trigger rate, ...





DETECTOR PERFORMANCE



Amaré et al., Eur. Phys. J. C (2019) 79:228, 1812.01472

Detector Response: duty cycle & stability

- Excellent **duty cycle**





- Good **total rate and gain stability**



Evolution of ^{109}Cd lines from calibrations along the whole data-taking (~ 2 year)

Detector response: threshold

Outstanding **light collection** of ~15 phe/keV

- Effectively **triggering below 1 keV$_{ee}$**



bulk ^{22}Na and ^{40}K events identified by coincidences with high energy gammas





Detector respose: threshold

Outstanding **light collection** of ~15 phe/keV

- Effectively **triggering below 1 keV$_{ee}$**



bulk ^{22}Na and ^{40}K events identified by coincidences with high energy gammas





- Energy threshold limited by PMT noise **filtering** protocols efficiency

- **Multiparametric cuts** to properly select events with pulse shapes from NaI(Tl) scintillation (efficiency computed on **^{109}Cd calibration and ^{22}Na and ^{40}K** coincidence populations)

Accepted event

REJECTED EVENTS



Blinded analysis

ANALYSIS STRATEGY

- Multiplicity-1 events in the RoI (1-6 keV) **blinded**
- We use multiplicity-2 events in the RoI and calibration events to tune the filtering algorithms and calculate the cut efficiencies
- We unblind 10% (~30 days randomly distributed along the first year) data for background assessment

10% unblinded data



Amaré et al., Eur. Phys. J. C (2019) 79:228, 1812.01472

Event selection & efficiency

CUTS

1. Pulse shape cut to select pulses with NaI(Tl) scintillation constant
2. We remove asymmetric events (<2 keVee) with origin in the PMT
3. Remove 1 s after a muon passage
4. Multiplicity = 1 (Reject events that deposit energy simultaneously in more than one crystal)



TOTAL EFFICIENCY



EFFICIENCY-CORRECTED BACKGROUND



Amaré et al., Eur. Phys. J. C (2019) 79:228, 1812.01

Background





ANAIS raw spectrum

ANAIS filtered spectrum

DAMA/LIBRA
Universe 4, 116 (2018), 1805.10486

Amaré et al., Eur. Phys. J. C (2019) 79:228, 1812.01472

Background model

At very low energy (<20 keV), main contamination in the crystal itself

- **^{40}K** and **^{22}Na** ($T_{1/2}$ = 2.6 y) peaks
- **^{210}Pb** (bulk+surface) ($T_{1/2}$ = 22.3 y)
- **^{3}H** ($T_{1/2}$ = 12.3 y)

Very good agreement with data except between 1-2 keV



Cosmogenic isotopes (^{3}H, ^{22}Na, …) and ^{210}Pb are decaying
→ prediction of the time depencence of the rate in the RoI



Amaré et al., Eur. Phys. J. C (2019) 79:412, 1812.01377

RESULTS ON ANNUAL MODULATION



Analysis strategy



Background decay rate in the RoI consistent with our background model (in green)

ANALYSIS STRATEGY

- Focus on model independent analysis searching from modulation
- In order to better compare with DAMA/LIBRA results, we use the **same energy regions ([1-6] keV, [2-6] keV) and fit parameters**
- Least square fit to

$$R(t) = R_0 + R_1 \exp(-t/\tau) + \boldsymbol{S_m} \cos(\omega(t+\phi))$$

Fixed parameters:

τ (background model)
ω (freq. corresponding to a period of 1 year)
ϕ (maximum in June, 2nd)

1st Annual modulation results: 1.5 y

157.55 kg x yr

Phys. Rev. Lett., 123, 031301 (2019)

Least squared fit to : $R(t) = R_0 + R_1 \exp(-t/\tau) + \boldsymbol{S_m} \cos(\omega(t+\phi))$

Fixed parameters:

τ (background model)

ω (freq. corresponding to a period of 1 year)

ϕ (maximum in June, 2nd)

$\boldsymbol{S_m}$ fixed to 0 in the null hypothesis and left unconstrained for the modulation hypothesis



DAMA/LIBRA result with 1 –free parameter is shown for comparison

1st Annual modulation results

157.55 kg x yr

[2-6] keV → S_m = −0.0044 ± 0.0058 c/keV/kg/d (S_m^{DAMA} = 0.0102 cpd/kg/keV)

[1-6] keV → S_m = −0.0015 ± 0.0063 c/keV/kg/d (S_m^{DAMA} = 0.0105 cpd/kg/keV)

- Null hypothesis is well supported by the χ^2test (p-values=0.18, 0.67)
- Best fits for the modulation hypothesis have p-values slightly lower than for the null hypothesis
- Best fits are compatible with no modulation and incompatible at 2.5σ (2-6 keV) and 1.9σ (1-6 keV) with DAMA/LIBRA results. Sensitivity (1.5 y) 1.8σ



The statistical significance of our result is determined by the standard deviation of the modulation amplitude distribution, $\sigma(S_m)$

arXiv:1903.03973

Phys. Rev. Lett., 123, 031301 (2019)

2 years results

213.6 kg x yr

M. L. Sarsa @ TAUP2019
J., Phys (Conference Series) 1468 (2020) 012014



2 years results

213.6 kg x yr

M. L. Sarsa @ TAUP2019

[2-6] keV → S_m = −0.0029 ± 0.0050 c/keV/kg/d (S_m^{DAMA} = 0.0102 cpd/kg/keV)

[1-6] keV → S_m = −0.0036 ± 0.0054 c/keV/kg/d (S_m^{DAMA} = 0.0105 cpd/kg/keV)

- Null hypothesis is well supported by the χ^2test (p-values=0.09, 0.63)
- Best fits for the mod. hypothesis p-values slightly lower than for the null hypothesis
- Best fits are compatible with no modulation and incompatible at 2.6σ with DAMA/LIBRA results. Present sensitivity 2σ

1.5 years (PRL 123 (2019) 031301)



2 years (TAUP 2019)



Sm & phase free

213.6 kg x yr

M. L. Sarsa @ TAUP2019

$$R(t) = R_0 + R_1 \exp(-t/\tau) + \boldsymbol{S_m} \cos(\omega(t + \phi))$$

Fixed parameters:

τ (background model)

ω (freq. corresponding to a period of 1 year)

Free parameters:

R_0, R_1, S_m, ϕ



3 years results

313.6 kg x yr

PRELIMINARY

NEW!



- **Preliminary sensitivity @ 3 years: 2.6σ**



3 year results

NEW!

313.6 kg x yr



ANAIS-112 SENSITIVITY





Calculating the sensitivity

Least squared fit to : $R(t) = R_0 + R_1 \exp(-t/\tau) + \boldsymbol{S_m} \cos(\omega(t + \phi))$

Three free parameters (R_0, R_1, S_m)



The experimental sensitivity is given by the standard deviation of the modulation amplitude $\boldsymbol{\sigma(S_m)}$, that can be calculated analytically from :

- Updated background
- Efficiency estimate and its error
- Live time distribution

See details in **Coarasa et al., Eur. Phys. J. C (2019) 79:233, 1812.02000**

Expected sensitivity

We quote our sensitivity to DAMA/LIBRA result as the ratio $S_m^{DAMA}/\sigma(S_m)$



Experimental sensitivity

We quote our sensitivity to DAMA/LIBRA result as the ratio $S_m^{DAMA}/\sigma(S_m)$



- **data confirm our sensitivity projection to DAMA/LIBRA result**
- **Sensitivity @ 2 years: 2σ**
- **Preliminary sensitivity @ 3 years: 2.6σ**

3σ sensitivity at reach in about 1 year from now!

Model dependent (SI interaction)

Likelihood 90% - 90%



- Standard halo model
- Spin-independent interaction
- $\rho_0 = 0.3\ \mathrm{GeV/cm^3}$
- $v_0 = 220\ \mathrm{km/s}$
- $v_{esc} = 650\ \mathrm{km/s}$
- $Q_{Na} = 0.30, Q_I = 0.09$

DAMA regions from:

C. Savage et al., JCAP04 (2009) 010

Summary

- Annual modulation is a distinctive signature of Dark Matter
- One positive signal (DAMA/LIBRA) for more than 20 years, in strong tension with other experiments
- Currently, many efforts trying to confirm / rule out DAMA/LIBRA signal with the same target. COSINE-100 and ANAIS-112 in data-taking
- ANAIS-112: is taking data in stable condition @ LSC since 3rd August 2017 with excellent performances. Up to now it has accumulated more than 300 kg×y exposure.
- **ANAIS-112 results** are compatible with absence of modulation and **incompatible with DAMA/LIBRA at 2σ after 2 years of data-taking**. Preliminary results for 3 years reach 2.6 sensitivity in [2-6] keV. **3σ sensitivity at reach in about 1 year from now!**

Summary

- Annual modulation is a distinctive signature of Dark Matter
- One positive signal (DAMA/LIBRA) for more than 20 years, in strong tension with other experiments
- Currently, many efforts trying to confirm / rule out DAMA/LIBRA signal with the same target. COSINE-100 and ANAIS-112 in data-taking
- ANAIS-112: is taking data in stable condition @ LSC since 3rd August 2017 with excellent performances. Up to now it has accumulated more than 300 kg×y exposure.
- **ANAIS-112 results** are compatible with absence of modulation and **incompatible with DAMA/LIBRA at 2σ after 2 years of data-taking**. Preliminary results for 3 years reach 2.6 sensitivity in [2-6] keV. **3σ sensitivity at reach in about 1 year from now!**

¡GRACIAS!